

T.G. Section III-A-2 Alternative Conservation System Part 2
[Only for FSA Compliance and if not Sodbusted]
Lordsburg Field Office
Irrigated Cropland Guide Sheet
Resource Data

MLRA 41 and 42

Soils: All irrigated soils in WEG 2 thru 7.

WEQ values: C-120 or less, I-134 or less, and L-3000 or less

If WEQ values listed above are exceeded, erosion losses for each rotation will be computed individually to ensure that total average wind erosion loss is within acceptable levels.

The following alternatives are acceptable regardless of the tillage method used provided the crop residues and/or growing crops are managed as indicated in the Management Requirements section to provide wind erosion protection during February to May, which is the critical blow season.

MANAGEMENT REQUIREMENTS

The Crop Rotation Needed for erosion control may include any combination of crops grown in any sequence that will significantly reduce erosion compared to his present system.

The existing Irrigation System will be maintained as irrigation is essential for crop production to control wind erosion. Land will be considered irrigated if adequate residues are produced to control wind erosion.

Alternative 1: Maintain Residues until Tilled Or Planted: The crop residue management for erosion protection requires leaving the residues from the previous crop on the surface until tillage operations for the next crop begins. After this, no residue is required if the land is planted, plowed or listed when irrigated. Avoid leaving the land in a smooth, dry, pulverized condition during the critical blow period.

Alternative 2: Growing crops: Maintain a 2 inch stubble on growing crops as alfalfa and small grain.

Alternative 3: Protect idle land with residues: If land is left fallow or idle manage the rotation where the idle land is preceded by a high residue, crop which has adequate residue for erosion protection. These residues will be maintained on the soil surface to leave the following "Small Grain Equivalents", (SGe) for the following soil types and Wind Erodibility Groups, (WEG). Loamy sands WEG-2, 2000# SGe/ac.; sandy loams, clays and highly calcareous loams in WEG-3, 4 and 4L, 1500# SGe/ac.; and loams in WEG-5, 6, and 7, 1250# SGe/ac.

The following crops normally produce the following SGe under normal management if left standing. Alfalfa, small grain, corn, milo, and chile produce over 3000# SGe/ac. Cotton produces about 2500# SGe/ac. If flat or shredded, chile will produce about 1600# SGe/ac. and cotton will produce less than 1500# SGe/ac.

Alternative 4: Idle land without residue: If inadequate residue is present and where adequate moisture is present on soils that will produce stable clods; plowing or listing is an adequate temporary alternative but should not exceed one year in the rotation. Irrigation may also be used.

Alternative 5: Idle land for extended periods: If land is to be left idle for extended periods, irrigation may be needed to re-establish the minimum residue, or the land may be planted to a perennial cover.

The Alternative Conservation System meets the erosion protection requirements of the Food Security Act of 1985 and is attainable and feasible within the Lordsburg Field Office.

Hidalgo SWCD Charles E. Sipe Date 6-13-88

District Conservationist James S. Wilber Date 6-13-88

Area Conservationist Stan Bublbaum, Acting Date 6/16/88

State Office Lay Margoz Date 7/7/88